

**NTP Technical Report
on Comparative Toxicity Studies of
o-, *m*-, and *p*-Chloroaniline**

(CAS Nos. 95-51-2, 108-42-9, and 106-47-8)

**Administered by Gavage
to F344/N Rats and B6C3F₁ Mice**

**Rajendra S. Chhabra, Ph.D., Study Scientist
National Toxicology Program
Post Office Box 12233
Research Triangle Park, NC 27709**

**United States Department of Health and Human Services
Public Health Service
National Institutes of Health**

Note to the Reader

The National Toxicology Program (NTP) is made up of four charter agencies of the United States Department of Health and Human Services (DHHS):

- the National Cancer Institute (NCI) of the National Institutes of Health;
- the National Institute of Environmental Health Sciences (NIEHS) of the National Institutes of Health;
- the National Center for Toxicological Research (NCTR) of the Food and Drug Administration; and
- the National Institute for Occupational Safety and Health (NIOSH) of the Centers for Disease Control.

In July 1981, the Carcinogenesis Bioassay Testing Program was transferred from NCI to NIEHS. NTP coordinates the relevant Public Health Service programs, staff, and resources that are concerned with basic and applied research and with biological assay development and validation.

The NTP develops, evaluates, and disseminates scientific information about potentially toxic and hazardous chemicals. This knowledge is used for protecting the health of the American people and for the primary prevention of disease.

The NTP designs and conducts studies to characterize and evaluate the toxicologic potential of selected chemicals in laboratory animals (usually two species, rats and mice). Chemicals selected for NTP toxicology studies are chosen primarily on the bases of human exposure, level of production, and chemical structure. The interpretive conclusions presented in this Toxicity Study Report are based only on the results of these NTP studies. Extrapolation of these results to other species and quantitative risk analyses for humans require wider analyses beyond the purview of these studies. Selection *per se* is not an indicator of a chemical's toxic potential.

The studies described in this Toxicity Study Report were performed under the direction of NIEHS and were conducted in compliance with NTP laboratory health and safety requirements. These studies met or exceeded all applicable federal, state, and local health and safety regulations. Animal care and use were in accord and compliance with the Public Health Service Policy on Humane Care and Use of Animals.

Listings of all published NTP reports and ongoing studies are available from NTP Central Data Management, NIEHS, P.O. Box 12233, MDE1-02, Research Triangle Park, NC 27709 (919-541-3419). Other information about NTP studies is available at the NTP's World Wide Web site:
<http://ntp-server.niehs.nih.gov>.

**NTP Technical Report
on Comparative Toxicity Studies of
o-, *m*-, and *p*-Chloroaniline**

(CAS Nos. 95-51-2, 108-42-9, and 106-47-8)

**Administered by Gavage
to F344/N Rats and B6C3F₁ Mice**

**Rajendra S. Chhabra, Ph.D., Study Scientist
National Toxicology Program
Post Office Box 12233
Research Triangle Park, NC 27709**

**United States Department of Health and Human Services
Public Health Service
National Institutes of Health**

CONTRIBUTORS

National Toxicology Program

Evaluated and interpreted results and reported findings

R.S. Chhabra, Ph.D., Study Scientist
J.R. Bucher, Ph.D.
L.T. Burka, Ph.D.
R.E. Chapin, Ph.D.
M.R. Elwell, D.V.M., Ph.D.
J. Mahler, D.V.M.
G.S. Travlos, D.V.M.
K.L. Witt, M.S., Integrated Laboratory Systems

Battelle Columbus Laboratories

Conducted studies and evaluated pathology findings

P.J. Kurtz, Ph.D., Principal Investigator
M.R. Hejtmancik, Ph.D.
R.L. Persing, D.V.M.
M.J. Ryan, D.V.M., Ph.D.
B.A. Trela, Ph.D.
J.T. Yarrington, D.V.M., Ph.D.

NTP Pathology Working Group

*o-Chloroaniline: Evaluated slides and prepared pathology report
(27 April 1993)*

P.K. Hildebrandt, D.V.M., Chairperson
PATHCO, Inc.
M.R. Elwell, D.V.M., Ph.D.
National Toxicology Program
W.F. MacKenzie, D.V.M., M.S.
Consultant
R.A. Herbert, D.V.M., Ph.D.
National Toxicology Program
R.C. Sills, D.V.M., Ph.D.
National Toxicology Program
S.A. Stefanski, D.V.M., M.S.
North Carolina State University

NTP Pathology Working Group

*m-Chloroaniline: Evaluated slides and prepared pathology report
(4 October 1994)*

P.K. Hildebrandt, D.V.M., Chairperson
PATHCO, Inc.
R. Cattley, V.M.D., Ph.D.
Chemical Industry Institute of Toxicology
F. Geoly, D.V.M., Observer
North Carolina State University
D.S. Marsman, D.V.M., Ph.D.
National Toxicology Program
A. Radovsky, D.V.M., Ph.D.
National Toxicology Program
C.C. Shackelford, D.V.M., M.S., Ph.D.
Experimental Pathology Laboratories, Inc.
R.C. Sills, D.V.M., Ph.D.
National Toxicology Program

Experimental Pathology Laboratories, Inc.

Provided pathology quality assurance

J. Peckham, D.V.M., M.S., Ph.D.
C.C. Shackelford, D.V.M., M.S., Ph.D.

Environmental Health Research and Testing, Inc.

Provided sperm motility and vaginal cytology evaluations

T. Cocanougher, B.A.
D.K. Gulati, Ph.D.
S. Russell, B.A.

Analytical Sciences, Inc.

Provided statistical analyses

R.W. Morris, M.S., Principal Investigator
S. R. Lloyd, M.S.
N.G. Mintz, B.S.

Biotechnical Services, Inc.

Prepared Toxicity Study Report

S.R. Gunnels, M.A., Principal Investigator
A.M. Macri-Hanson, M.A., M.F.A.
W.D. Sharp, B.A., B.S.

PEER REVIEW

The draft report on the toxicity studies of *o*-chloroaniline, *m*-chloroaniline, and *p*-chloroaniline was evaluated by the reviewers listed below. These reviewers serve as independent scientists, not as representatives of any institution, company, or governmental agency. In this capacity, reviewers determine if the design and conditions of these NTP studies are appropriate and ensure that the toxicity study report presents the experimental results and conclusions fully and clearly.

Gary P. Carlson, Ph.D.
School of Health Sciences
Purdue University
West Lafayette, IN


John Mennear, Ph.D.
Raleigh, NC

CONTENTS


ABSTRACT	5
INTRODUCTION	7
Physical Properties, Production, Use, and Exposure	7
Disposition and Metabolism	8
Toxicity	8
Study Rationale and Design	12
MATERIALS AND METHODS	13
Procurement and Characterization of <i>o</i> -Chloroaniline and <i>m</i> -Chloroaniline	13
Dose Formulations	13
Toxicity Study Designs	14
Statistical Methods	19
Quality Assurance	19
Genetic Toxicity Studies	20
RESULTS	25
13-Week Gavage Studies in F344/N Rats	25
13-Week Gavage Studies in B6C3F ₁ Mice	46
Genetic Toxicity	62
DISCUSSION	65
REFERENCES	71
APPENDIXES	
Appendix A Summary of Nonneoplastic Lesions in Rats	A-1
Appendix B Summary of Nonneoplastic Lesions in Mice	B-1
Appendix C Organ Weights and Organ-Weight-to-Body-Weight Ratios	C-1
Appendix D Hematology and Clinical Chemistry Results	D-1
Appendix E Reproductive Tissue Evaluations and Estrous Cycle Characterization	E-1
Appendix F Genetic Toxicology	F-1

ABSTRACT


***o*-Chloroaniline**



***m*-Chloroaniline**



***p*-Chloroaniline**



CAS Number 95-51-2
Synonyms *o*-aminochlorobenzene
 2-chlorobenzenamine
 2-chloroaniline
 2-chlorophenylamine

108-42-9
m-aminochlorobenzene
 3-chlorobenzenamine
 3-chloroaniline
 3-chlorophenylamine

106-47-8
p-aminochlorobenzene
 4-chlorobenzenamine
 4-chloroaniline
 4-chlorophenylamine

Molecular Formula C₆H₆ClN
Molecular Weight 127.57

Chlorinated anilines are used as intermediates in the manufacture of dyes, drugs, and agricultural agents. In comparative 13-week studies conducted to determine the structure-toxicity relationships of *o*-, *m*-, and *p*-chloroaniline, groups of 10 male and 10 female F344/N rats and B6C3F₁ mice were administered 0, 10, 20, 40, 80, or 160 mg *o*- or *m*-chloroaniline per kilogram body weight in dilute hydrochloric acid by gavage. Animals were evaluated for hematology, clinical chemistry, histopathology, and reproductive system effects. Genetic toxicity studies of *o*- and *m*-chloroaniline *in vivo* and *in vitro* were also conducted. The results of the *o*- and *m*-chloroaniline studies were compared to results from the *p*-chloroaniline studies performed previously under similar experimental conditions by the same laboratory; doses in the *p*-chloroaniline studies were 0, 5, 10, 20, 40, and 80 mg/kg for rats and 0, 7.5, 15, 30, 60, and 120 mg/kg for mice.

The hematopoietic system was the target of *o*-, *m*-, and *p*-chloroaniline in rats and mice. Neither the *o*- nor the *p*- isomer had an adverse effect on survival; the death of one female rat in the 160 mg/kg *m*-chloroaniline group during week 12 was possibly secondary to methemoglobinemia. The final mean body weights and weight gains of male rats in the highest dose group in each study and female mice in the 160 mg/kg group in the *o*-chloroaniline study were significantly less than those of the respective controls. Clinical findings of toxicity included a transient bluish discoloration of the genital and footpad regions in rats administered *o*- or *m*-chloroaniline and tremors in rats and mice administered *o*-chloroaniline and in mice administered

m-chloroaniline; these effects occurred primarily in the 80 and 160 mg/kg groups. Methemoglobin concentrations were increased in dosed rats and mice in all studies and resulted in a secondary anemia; the severity of the anemia increased with increasing dose. Microscopic lesions considered related to chemical administration in rats and mice included hemosiderin pigmentation in the bone marrow, kidney, liver, and spleen; hematopoiesis in the liver and spleen; and erythroid cell hyperplasia in the bone marrow. These lesions reflected the response to hemolytic anemia and methemoglobinemia induced by the chloroanilines. A comparative analysis of the results suggests that *p*-chloroaniline is the most potent of the chloroaniline isomers in the induction of methemoglobin formation in rats and mice, followed by *m*-chloroaniline and then by *o*-chloroaniline. This order of potency was also observed for changes in other hematology parameters and in spleen weights, gross and microscopic lesions, and the severity of hemosiderin deposition.

Although the *o*-, *m*-, and *p*- isomers of chloroaniline all exhibit genetic toxicity, the profiles of activity among the three isomers are not identical. *p*-Chloroaniline was mutagenic in all assays in which it was tested, including the *Salmonella* assay, the mouse lymphoma assay, *in vitro* Chinese hamster ovary cell cytogenetics assays, and the *in vivo* mouse bone marrow micronucleus assay; in contrast, *o*- and *m*-chloroaniline gave mixed results among the various assays in which each was tested.

In conclusion, chloroaniline isomers are hematotoxic and have the same pattern of toxicity in rats and mice. Hematotoxicity occurred at all doses in these studies. *p*-Chloroaniline induces the most severe hematotoxic effect, followed by *m*-chloroaniline, then *o*-chloroaniline. Each of the three isomers is more toxic to rats than to mice. *p*-Chloroaniline is clearly genotoxic in various test systems, while the results for the *o*- and *m*- isomers are inconsistent and indicate weak or no genotoxic effects.